

Recombinant Human MAF protein (His Tag)

Catalog Number: PDEH101086

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Human |
| Source | E.coli-derived Human MAF protein Met1-Met373, with an N-terminal His |
| Calculated MW | 40.9 kDa |
| Observed MW | 45 kDa |
| Accession | O75444-2 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95% as determined by reducing SDS-PAGE. |
| Endotoxin | < 10 EU/mg of the protein as determined by the LAL method |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol. |
| Reconstitution | It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

c-Maf is an approximately 40 kDa transcriptional regulator that contains one bZIP domain. It can associate into homodimers and heterodimers with other basic leucine zipper transcription factors. c-Maf plays an important role in fetal erythropoiesis, lens development, mechanosensory neuron development, and the differentiation of chondrocytes and osteoblasts. In the pancreas, c-Maf promotes the transcription of glucagon in alpha cells and insulin in beta cells.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.4