A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse Lumican/LUM Protein (His Tag)

Catalog Number: PKSM041313

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}		crip					
	00	0	РΤ	n	ŤΤ		m
v	\mathbf{c}	v.		w	w	w	ш

Species Mouse

Source HEK293 Cells-derived Mouse Lumican/LUM protein Gln 19-Asn338, with an C-terminal

His

Calculated MW 37.4 kDa
Observed MW 40-60 kDa
Accession P51885

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



Background

Keratan sulfate proteoglycan lumican, mostly known as lumican, belongs to the SLRP class II subfamily and the small leucine-rich proteoglycan (SLRP) family. Lumican binds to laminin. It has positive regulation of transcription from RNA polymerase II promoter and transforming growth factor beta1 production. Lumican is expressed in a variety of tissues, including skin, artery, lung, cornea, kidney, bone, aorta, and articular cartilage. The abnormal connective tissue phenotype seen in the Lumican null mice shows the importance of the role of Lumican in collagen fibrillogenesis. In addition to the control of collagen fibril assembly, Lumican has been shown to play a role in the regulation of cell proliferation, migration, and adhesion.

For Research Use Only