A Reliable Research Partner in Life Science and Medicine

Recombinant Human SMYD3/ZMYND1 Protein (GST Tag)

Catalog Number: PKSH031202

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source Baculovirus-Insect Cells-derived Human SMYD3/ZMYND1 protein Lys 35-Ser 369,

with an N-terminal GST

 Mol_Mass
 65.6 kDa

 Accession
 NP_073580.1

Bio-activity Not validated for activity

Properties

Purity > 88 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

ShippingThis product is provided as lyophilized powder which is shipped with ice packs. **Formulation**Lyophilized from sterile 20mM Tris, 150mM NaCl, 0.5mM DTT, 0.5mM GSH, pH 8.0


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 88 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Fax: 1-832-243-6017

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

SET and MYND domain-containing protein 3, also known as Zinc finger MYND domain-containing protein 1, SMYD3, and ZMYND, is a member of the histone-lysine methyltransferase family. SMYD3 contains oneMYND-type zinc finger and oneSET domain. SMYD3 is a histone H3 lysine-4-specific methyltransferase. It is expressed in skeletal muscles and testis. It is overexpressed in a majority of colorectal carcinoma (CRC) and hepatocellular carcinoma (HCC). SMYD3 plays an important role in transcriptional regulation in human carcinogenesis. It activates the transcription of a set of downstream genes. Of these downstream genes, there are several oncogenes and genes associated with cell adhesion (including those of N-Myc, CrkL, Wnt10b, L-selectin, CD31 and galectin-4), which have been shown to have effects on cell viability, adhesion, migration and metastasis. Increased SMYD3 expression is essential for the proliferation of breast cancer cells. SMYD3 may be a promising new target of therapeutic intervention for the treatment of cancers or other pathological processes associated with cell adhesion and migration.

Fax: 1-832-243-6017