Elabscience®

PE/Elab Fluor[®] 594 Anti-Rat CD3 Antibody[G4.18]


Catalog Number: E-AB-F1228UP

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Rat
Host	Mouse
lsotype	Mouse IgG3, κ
Clone No.	G4.18
Isotype Control	PE/Elab Fluor [®] 594 Mouse IgG3, κ Isotype Control[A112-3] [Product E-AB-F09752P]
Conjugation	PE/Elab Fluor [®] 594
Conjugation Information	PE/Elab Fluor [®] 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm
	(e.g., a 610/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].





Rat splenocytes are stained with PE/Elab Fluor[®] 594 Anti-Rat CD3 Antibody[G4.18] (filled gray histogram) or PE/Elab

Fluor $^{\mbox{\scriptsize R}}$ 594 Mouse IgG3, κ Isotype Control (empty black histogram).

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged
	exposure to light and do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	T-cell surface glycoprotein CD3 δ ; γ ; ϵ ;and ζ chains;CD3 Complex;T3;CD3
Uniprot ID	P19377;Q64159;D4A5M2
Gene ID	25710,300678,315609,25300

For Research Use Only

Elabscience®

.

Elabscience Biotechnology Co., Ltd. A Reliable Research Partner in Life Science and Medicine

Background

CD3 is a complex composed of δ , γ , ϵ , and ζ chains. They are 20-25 kD members of the immunoglobulin superfamily and associated with the T cell receptor (TCR). CD3 is expressed on thymocytes, peripheral T cells, some NK-T cells, and dendritic epidermal T cells. CD3 is involved in antigen recognition, signal transduction, and T cell activation