Recombinant Mouse PTPN2/PTPT Protein (aa 2-314, His Tag)

Catalog Number: PKSM040482

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	Baculovirus-Insect Cells-derived Mouse PTPN2/PTPT protein Ser 2-Asn 314, with an N-terminal His
Calculated MW	38.7 kDa
Observed MW	42 kDa
Accession	Q06180-1
Bio-activity	Measured by its ability to dephosphorylate a phosphotyrosine residue in an EGF receptor 988-998 phosphopeptide substrate, R&D Systems, Catalog # ES006. The specific activity is > 15 nmoles/min/μg.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 20% gly, 0.1mM TCEP
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Tyrosine-protein phosphatase non-receptor type 2, also known as T-cell protein-tyrosine phosphatase, PTPN2 and PTPT, is a cytoplasm protein which belongs to theprotein-tyrosine phosphatase family and Non-receptor class 1 subfamily. Members of the protein tyrosine phosphatase (PTP) family share a highly conserved catalytic motif, which is essential for the catalytic activity. TC-PTP / PTPN2 is a cytosolic tyrosine phosphatase that functions as a negative regulator of a variety of tyrosine kinases and other signaling proteins. The expression of TC-PTP / PTPN2 plays a role of tumor suppressor and may modulate response to treatment. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. Epidermal growth factor receptor and the adaptor protein Shc were reported to be substrates of this PTP, which suggested the roles in growth factor mediated cell signaling. TC-PTP / PTPN2 is an enzyme that is essential for the proper functioning of the immune system and that participates in the control of cell proliferation, and inflammation.TC-PTP / PTPN2 was identified as a negative regulator of NUP214-ABL1 kinase activity.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com