

Recombinant Mouse Frizzled-10/FZD10 Protein (Fc Tag)

Catalog Number: PKSM040610

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse Frizzled-10/FZD10 protein Met 1-Gly 162, with an C-terminal hFc
Calculated MW	43.0 kDa
Observed MW	52 kDa
Accession	NP_780493.1
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Frizzled-10, also known as Fz-10, CD350 and FZD10, is a multi-pass membrane protein which belongs to the G-protein coupled receptor Fz/Smo family. Frizzled-10 / FZD10 is abundantly expressed in the cerebellum, followed by cerebral cortex, medulla and spinal cord; very low levels in total brain, frontal lobe, temporal lobe and putamen. It is weakly expressed in adult brain, heart, lung, skeletal muscle, pancreas, spleen and prostate. Frizzled-10 / FZD10 is a receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. Frizzled-10 / FZD10 may also be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017