Recombinant Human LILRB4/CD85k/ILT3 Protein (aa 22-259, Fc Tag)



Catalog Number: PKSH033604

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Leukocyte immunoglobulin-like receptor subfamily B member 4;Mast cell surface

Lyophilized from a 0.2 µm filtered solution of PBS;pH 7.4.glycoprotein

Gp49B;CD85k;Lilrb4;Gp49b

Species Human

Expression Host HEK293 Cells
Sequence Gln22-Glu259
Accession Q8NHJ6
Calculated Molecular Weight 53.2 kDa
Observed molecular weight 65-85 kDa
Tag C-Fc

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Mouse Leukocyte Immunoglobulin-like Receptor Subfamily B Member 4 (LILRB4/CD85k/ILT3) is an approximately transmembrane glycoprotein that negatively regulates immune cell activation. Mouse LILRB4 consists of a 215 amino acid (aa) extracellular domain with two Ig-like domains; a 22 aa transmembrane segment; and a 75 aa cytoplasmic domain with 3 immunoreceptor tyrosine-based inhibitory motifs (ITIM). Within the ECD; mouse LILRB4 shares 45% and 77% aa sequence identity with human and rat LILRB4; respectively. Alternative splicing of mouse LILRB4 generates a potentially soluble isoform that lacks the transmembrane segment. LILRB4 is expressed on dendritic cells (DC);

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human LILRB4/CD85k/ILT3 Protein (aa 22-259, Fc Tag)



Fax: 1-832-243-6017

Catalog Number: PKSH033604

monocytes; macrophages; and vascular endothelial cells (EC). Ligation of LILRB4 triggers ITIM-mediated inhibition of cellactivating signaling; leading to enhanced immune tolerance and reduced allogeneic graft rejection. Soluble LILRB4 induces the differentiation of CD8+ T suppressor cells (Ts) that can inhibit the effector functions of CD4+ Th cells and CD8+ CTL. In turn; CD8+ Ts cells induce LILRB4 up-regulation and a tolerogenic phenotype in monocytes; DC; and EC.

Web: www.elabscience.com Email: techsupport@elabscience.com