## **Elabscience**®

## **QTRT1** Polyclonal Antibody

## catalog number: E-AB-52610

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human QTRT1
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000
Data	
Western blot analysis of Polyclonal Antil <b>Observed-M</b>	<ul> <li><sup>55-</sup></li> <li><sup>36-</sup></li> <li><sup>28-</sup></li> <li><sup>17-</sup></li> <li><sup>17-</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>17</sup></li> <li><sup>18</sup></li> <li><sup>18</sup></li> <li><sup>18</sup></li> <li><sup>18</sup></li> <li><sup>19</sup></li> <li><sup>1</sup></li></ul>
Preparation & Storage	
r reparation & Storage	
Storago	Store at _20°C Valid for 12 months Avoid freeze / they cycles
Storage Shipping	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack, upon receipt, store it immediately at the

This gene encodes the catalytic subunit of tRNA-guanine transglycosylase. tRNA-guanine transglycosylase is a heterodimeric enzyme complex that plays a critical role in tRNA modification by synthesizing the 7-deazaguanosine queuosine, which is found in tRNAs that code for asparagine, aspartic acid, histidine and tyrosine. A pseudogene of this gene is located on the long arm of chromosome X.