

# FITC-conjugated Myc-Tag Monoclonal Antibody

catalog number: AN00300C

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	All
<b>Immunogen</b>	MYC tag fusion protein
<b>Host</b>	Mouse
<b>Isotype</b>	IgG1
<b>Clone</b>	6B10
<b>Purification</b>	Protein A/G Purification
<b>Conjugation</b>	FITC(Ex/Em:495nm/525nm)
<b>buffer</b>	PBS with 0.05% proclin 300 and 0.1% gelatin,pH7.4

## Applications

## Recommended Dilution

<b>FCM</b>	0.2 µg per 10 <sup>6</sup> cells in a 100 µL suspension
------------	---

## Data



1X10<sup>6</sup> HEK-293F cells were intracellularly stained with 0.2 µg FITC-conjugated Myc-Tag Monoclonal Antibody (red), or 0.2 µg FITC-conjugated Myc-Tag Monoclonal Antibody Isotype Control (green).

## Preparation & Storage

<b>Storage</b>	Store at 4°C Valid for 12 months. Avoid freeze / thaw cycles. Protected from prolonged exposure to light.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

The MYC tag, derived from the c-MYC protein, is a popular epitope tag for detecting the expression of recombinant proteins in yeast, bacteria, insect, and mammalian cell systems. It can be used for affinity chromatography, then used to separate recombinant

## For Research Use Only