A Reliable Research Partner in Life Science and Medicine

Recombinant Human EphB2 Protein (Fc Tag)

Catalog Number: PKSH032012

Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip					
	00	CI	411	m	П	n	m
JU	\mathbf{c}	U		J.	"	v.	ш

Species Human

Source HEK293 Cells-derived Human EphB2 protein Val19-Ser482, with an C-terminal Fc

 Mol_Mass
 78.5 kDa

 Accession
 Q6NVW1

Bio-activity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Ephrin type-B receptor 2(EPHB2) belongs to the protein kinase superfamily and Ephrin receptor subfamily. EPHB2 contains 1 Eph LBD domain; 2 fibronectin type-III domains; 1 protein kinase domain and 1 SAM domain. Ephrin receptors and their ligands; the ephrins; mediate numerous developmental processes; particularly in the nervous system. Based on their structures and sequence relationships; ephrins are divided into the ephrin-A (EFNA) class; which are anchored to the membrane by a glycosylphosphatidylinositol linkage; and the ephrin-B (EFNB) class; which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family.

For Research Use Only