Elabscience®

FLT4 Polyclonal Antibody

catalog number: E-AB-53195

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human	
Immunogen	Synthetic peptide of human FLT4	
Host	Rabbit	
Is otype	IgG	
Purification	Antigen affinity purification	
Conjugation	Unconjugated	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilution	
IF	1:100-1:500	
Data		
5. T.		12 m
Immunofluorescence a	analysis of hela cell using FLT4	Immunofluorescence analysIs of HepG2 cell using FLT4
	analysis of hela cell using FLT4 ibody at dilution of 1:100	Immunofluorescence analysIs of HepG2 cell using FLT4 Polyclonal Antibody at dilution of 1:100
Polyclonal Anti	ibody at dilution of 1:100	
Polyclonal Anti Preparation & Storage	ibody at dilution of 1:100 Store at -20°C Valid for 1	Polyclonal Antibody at dilution of 1:100

This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA. Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFC and VEGFD, and plays an essential role in adult lymphangiogenesis and in the development of the vascular network and the cardiovascular system during embryonic development. Promotes proliferation, survival and migration of endothelial cell s, and regulates angiogenic sprouting.