Purified Anti-Human CD340 Antibody[chA21]

catalog number: AN003520P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						٠,		
D	Α		C	PТ	m	ŤΤ	U.	m
\mathbf{L}	v	0	v.	ш	עו	UΙ	U.	

Reactivity Human

Immunogen Recombinant Human CD340 protein

Host Mouse

Isotype Mouse IgG2a, κ

Clone chA21

Purification >98%, Protein A/G purified

Conjugation Unconjugated


buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze

to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$

Data



MCF-7 were stained with 0.2 μg Purified Anti-Human CD340 Antibody[chA21] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left), followed by FITC-conjugated Goat

Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

ErbB2, also called Neu and Her2, is a transmembrane glycoprotein in the ErbB family of tyrosine kinase receptors for EGF superfamily growth factors. ErbB2 is widely expressed in epithelial cells and over-expressed in a large number of breast carcinomas. ErbB2 has no identified ligands but heterodimerizes with ErbB1/EGF R, ErbB3, or ErbB4 to form higher affinity signaling complexes. The protease ADAM10 releases a 110 kDa soluble fragment of ErbB2 from the cell surface. ErbB2 plays roles in development, cancer, communication at the neuromuscular junction, and regulation of cell growth and differentiation. The ErbB2/ErbB3 heterodimer is expressed in the majority of breast, skin, ovary and gastrointestinal tumors and transduces a highly mitogenic signal in response to neuregulin 1 (NRG1; heuregulin 1) or NRG2. ErbB3, ErbB2 and neuregulin are all required for formation of the sympathetic nervous system.

For Research Use Only