Recombinant Human CDK5 Protein (GST Tag)

Catalog Number: PKSH031366

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human CDK5 protein Met 1-Pro 292, with an N-terminal GST
Calculated MW	59.6 kDa
Observed MW	59.6 kDa
Accession	NP_004926.1
Bio-activity	Not validated for activity

Properties

Purity	> 94% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, 0.5 mM GSH, pH 7.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 94 % as determined by reducing SDS-PAGE.

Background

Cell division protein kinase 5, also known as Cyclin-dependent kinase 5, Serine/threonine-protein kinase PSSALRE, Tau protein kinase II catalytic subunit, TPKII catalytic subunit and CDK5, is a cytoplasm protein which belongs to theprotein kinase superfamily, CMGC Ser/Thr protein kinase family and CDC2/CDKX subfamily. Cyclin-dependent kinases (Cdks) are a family of proline-directed Ser/Thr kinases known for their role in the control of cell cycle progression. In 1992, this family was joined by CDK5, which is an atypical member in that it uses its own activators and is multifunctional, playing important regulatory roles in multiple cellular functions. CDK5, unlike other Cdks, is not regulated by cyclins, and its activity is primarily detected in postmitotic neurons in developing and adult nervous systems. CDK5 is activated by association with a neuron-specific activator, p35 or its isoform p39. CDK5 is probably involved in the control of the cell cycle. It interacts with D1 and D3-type G1 cyclins. CDK5 can phosphorylate histone H 1, tau, MAP2 and NF-H and NF-M. It also interacts with p35 which activates the kinase. CDK5 plays important roles in various neuronal activities, including neuronal migration, synaptic activity, and neuronal cell death.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com