

**TGFβ 1/TGFB1, Swine, Recombinant****Cat. No. : GPCK346****产品信息**

物种	Swine
表达宿主	E.coli
序列信息	MALDTNYCFSSSTEKNCCVRQLYIDFRKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKV LALYNQHNP GASAAPCCVPQALEPLPIVYYVGRKPKVEQLSNMIVRSCKCS with polyhistidine tag at the C-terminus.
检索号	P07200.2
标签	His-tag at the C-terminus
分子量	13.7 kDa
有效期	12 months
生物活性	Measure by its ability to inhibit IL-4-induce proliferation in HT-2 cells. The ED50 for this effect is < 0.1 ng/mL.

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 20 mM sodium citrate, 0.2 M NaCl, pH 4.5.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein. In some experiments, it recommends to add 10 mM HCl when reconstitute lyophilized protein.

**背景介绍**

Transforming growth factor beta 1 or TGF-β1 is a polypeptide member of the transforming growth factor beta superfamily of cytokines. It is a secreted protein that performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation, and apoptosis. In humans, TGF-β1 is encoded by the TGFB1 gene.

