A Reliable Research Partner in Life Science and Medicine

PE Anti-Mouse CD115/CSF-1R Antibody[AFS98]

Catalog Number: E-AB-F1107D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ
Clone No. AFS98

Isotype Control PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



C57BL/6 murine abdominal macrophages are stained with APC Anti-Mouse CD11b Antibody and PE Anti-Mouse CD115 Antibody (Left). Abdominal macrophages are stained with APC Anti-Mouse CD11b Antibody and PE Rat IgG2a, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD115;CSF-1 receptor (EC:2.7.10.1);CSF-1-R;CSF-1R;Csf1r;Csfmr;Fms;M-CSF-R;

Macrophage colony-stimulating factor 1 receptor; Proto-oncogene c-Fms

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Uniprot ID P09581
Gene ID 12978
Background CSF-1R

CSF-1R, also known as CD115 and M-CSFR, is a single-pass type I membrane protein and member of the platelet-derived growth factor receptor family. This c-fms (Fms proto-oncogene) gene product's natural ligands include M-CSF and IL-34. Structural studies of CD115 have described an Ig-like extracellular domain, a transmembrane domain, an intracellular juxtamembrane domain, a split tyrosine kinase domain, and a C-terminal tail receptor. Receptor activation induces homodimerization in addition to phosphorylation and ubiquitination of intracellular residues. CD115 directly influences tissue macrophage and osteoclast differentiation and proliferation. It is expressed on monocytes/macrophages, peritoneal exudate cells, plasmacytoid and conventional dendritic cells, and osteoclasts.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com