Recombinant Human PD-1/PDCD1 Protein (mFc Tag)

Catalog Number: PKSH032864



Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}					٠,		
H)	es	C1	ľ٦	n	П	on	

 Species
 Human

 Mol_Mass
 42.0 kDa

 Accession
 Q15116

Bio-activity Immobilized Anti-Human PD-1 mAb at 2µg/ml (100 µl/well) can bind Human PD-1-

mFc. The ED50 of Human PD-1-mFc is 12.53 ng/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU} \text{ per } \mu\text{g} \text{ of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

PD-1 is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 inhibits the T-cell proliferation and production of related cytokines including IL-1; IL-4; IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition; coligation of PDCD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. PDCD1 has been suggested to be involved in lymphocyte clonal selection and peripheral tolerance; and thus contributes to the prevention of autoimmune diseases. As a cell surface molecule; PDCD1 regulates the adaptive immune response. Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation; cytokine production; and cytolytic function.

For Research Use Only