

Recombinant TNF receptor II Monoclonal Antibody

catalog number: **AN301888L**


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human;Mouse
Immunogen	Recombinant human TNF receptor II fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A604
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB	1:500-1:1000
IHC	1:50-1:100
IF	1:50



Immunohistochemistry of paraffin-embedded Mouse liver using TNF receptor II Monoclonal Antibody at dilution of 1:100.



Immunofluorescent analysis of (4% Paraformaldehyde) fixed Jurkat cells using anti-TNF receptor II Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

TNF receptor II (TNF-R2) is one of the receptors of TNF- α . The two receptors for TNF- α , TNF-R1 (55 kDa) and TNF-R2 (75 kDa), can mediate distinct cellular responses. In most cases, cytotoxicity elicited by TNF has been reported to act through TNF-R1. In contrast, TNF-R2 appears to be important in T cell signaling and responses to infection. TNF-R2 binds to distinct members of the TRAF family, leading to the activation of NF- κ B. Soluble forms of both receptors have also been characterized which can bind TNF- α and may play an important role in immune disorders.

For Research Use Only