普诺赛[®] Procell system

ACVR2A (C-6His), Human, Recombinant

Cat. No.: GPCK106

产品信息

物种

Human Cells Cell system 表达宿主

Ala20-Pro134 序列信息

检索号 P27037 分子量 14.35 kDa 有效期 6 months

产品特性

内毒素 (EU/μg) < 0.1

保存 Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months

under sterile conditions after opening. Please minimize freeze-thaw cycles.

运输 The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the

temperature listed below.

制剂 Supplied as a 0.2 µm filtered solution of 20 mM PB, 300 mM NaCl, 10% Glycerol, 5%

Trealose, pH 7.4.

背景介绍

Activin Receptor Type-2A is a Protein that in humans is encoded by the ACVR2A gene. ACVR2A is an activintype 2 Receptor. This gene encodes activin A type II Receptor. Activins are dimeric growth and differentiation factors which belong to the transforming Growth Factor-beta (TGF-beta) superfamily of structurally related signaling Proteins. Activins signal through a heteromeric complex of Receptor serinekinases which include at least two type I (I and IB) and two type II (II and IIB) Receptors. These Receptors areall transmembrane Proteins, composed of a Ligand-binding extracellular domain with cysteine-rich region, atransmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type IReceptors are essential for signaling; and type II Receptors are required for binding Ligands and forexpression of type I Receptors. Type I and II Receptors form a stable complex after Ligand binding, resultingin phosphorylation of type I Receptors by type II Receptors. Type II Receptors are 普诺赛[®] | Procell system considered to beconstitutively active kinases.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



