

**FGF-3/HBGF-3/INT 2, Human, Recombinant****Cat. No. : GPCK293****产品信息**

物种	Human
表达宿主	E.coli
序列信息	MDAGGRGGVYEHLLGGAPRRRKLYCATKYHLQLHPSGRVNGSLENSAYSILEITAVEVGIVAIRG LFSGRYLAMNKRGRLYASEHYSACEFVERIHELGYNTYASRLYRTVSSTPGARRQPSAERLW YVSVNGKGRPRRGFKTRRTQKSSLFLPRVLDHRDHEMVRQLQSGLP RPPGKGVQPRRRR with polyhistidine tag at the C-terminus.
检索号	P11487.1
标签	His-tag at the C-terminus
分子量	21.99 kDa
有效期	12 months
生物活性	Measure by its ability to induce 3T3 cells proliferation. The ED50 for this effect is < 78 ng/mL.

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**背景介绍**

Fibroblast Growth Factor 3 (FGF-3) belongs to the large FGF family which has at least 23 members (1, 2). All FGF family members are heparin-binding growth factors with a core 120 amino acid (aa) FGF domain that allows for a common tertiary structure. Studies have suggested that FGF-3 and FGF-8 function synergistically in otic placode induction and during neuronal development to regulate dorsoventral axis formation. During development, the activities of FGF-3 and FGF-8 are regulated negatively by the sprouty family proteins and by Sef (similar expression t).

