

A Reliable Research Partner in Life Science and Medicine

Recombinant ABHD14B Monoclonal Antibody

catalog number: AN300273P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human ABHD14B protein

Host Rabbit Isotype lgG Clone B191 **Purification** Protein A


Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

1:500-1:1000 **WB**

ΙP 0.2-1 µL/mg of lysate

Data



Immunoprecipitation analysis using 0.5 µL anti-ABHD14B Western Blot with ABHD14B Monoclonal Antibody at dilution Monoclonal Antibody and 15 µl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate

using ABHD14B Monoclonal Antibody at a dilution of 1:500.

Lane A:0.5 mg Jurkat Whole Cell Lysate, Lane B:0.5 mg

Hela Whole Cell Lysate Observed-MW:25 kDa Calculated-MW:25 kDa of 1:500. Lane A: Jurkat Whole Cell Lysate,

Lysates/proteins at 30 µg per lane.

Observed-MW:25 kDa Calculated-MW:25 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

> activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

ABHD14B belongs to the AB hydrolase superfamily, ABHD14 family. It can be detected in spleen, thymus, prostate, testis, ovary, small intestine, colon, peripheral blood leukocyte, heart, placenta, lung, liver, skeletal muscle, pancreas and kidney. ABHD14B has hydrolase activity towards p-nitrophenyl butyrate (in vitro) and may interact with TAF1. It may activate transcription. Recombinant human ABHD14B protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques. ABHD14B contains an alpha/beta hydrolase fold, which is a catalytic domain found in a very wide range of enzymes. In molecular biology, the alpha/beta hydrolase fold is common to a number of hydrolytic enzymes of widely differing phylogenetic origin and catalytic function. The Ab hydrolase domain containing gene subfamily is comprised of 15 mostly uncharacterized members.

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Rev. V1.2