

## Purified Anti-Mouse CD19 Antibody[1D3], Functional Grade

catalog number: E-AB-F09860

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, κ
<b>Clone</b>	1D3
<b>Purification</b>	>98%, Protein A/G purified
<b>Buffer</b>	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

### Applications


### Recommended Dilution

<b>FCM</b>	≤ 0.2 µg per million cells in 100 µL volume
<b>Depletion</b>	Reported in the literature
<b>Neut</b>	Reported in the literature

### Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse CD19 Antibody[1D3], Functional Grade (Right) and 0.2 µg Rat IgG2a, κ Isotype Control (Left), followed by PE-conjugated Goat Anti-Rat IgG Secondary Antibody.



Monomer purity ≥102% as determined by analytical size-exclusion chromatography (SEC)

### Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
<b>Shipping</b>	Ice bag

### Background

CD19 is a 95 kD glycoprotein also known as B4. It is a member of the Ig superfamily, expressed on all pro-B to mature B cells (during development) and follicular dendritic cells. Plasma cells do not express CD19. CD19, in association with CD21 and CD81, forms a molecular complex integral to B cell activation.

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

### Application References

Min Dai, et al. Clin Cancer Res. 2015 Mar 1;21(5):1127-38. Yaron Carmi, et al. Nature. 2015 May 7;521(7550):99-104.

### For Research Use Only