# **PCBD1 Polyclonal Antibody**

catalog number: E-AB-19099



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human;Mouse;Rat  |
| Immunogen    | Fusion protein of human PCBD1  |
| Host         | Rabbit   |
| Is otype     | IgG  |
| Purification | Antigen affinity purification  |
| Conjugation  | Unconjugated   |
| buffer       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

| Applications | Recommended Dilution |
|--------------|----------------------|
| WB           | 1:500-1:2000         |
| IHC          | 1:50-1:300           |

### Data






cancer tissue using PCBD1 Polyclonal Antibody at dilution

of 1:60(×200)

Western blot analysis of Mouse kidney tissue lysate using

PCBD1 Polyclonal Antibody at dilution of 1:800



## Calculated-MV:12 kDa



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PCBD1 Polyclonal Antibody at dilution of 1:60(×200)

| Preparation & Storage |  |
|-----------------------|--|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.  |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

### Background

This gene encodes a member of the pterin-4-alpha-carbinolamine dehydratase family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The encoded protein functions as both a dehydratase involved in tetrahydrobiopterin biosynthesis, and as a cofactor for HNF1A-dependent transcription. A deficiency of this enzyme leads to hyperphenylalaninemia. Alternative splicing results in multiple transcript variants.

# For Research Use Only