

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Human CD122/IL-2RB Antibody[TU27]

Catalog Number: E-AB-F1151C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ

Clone No. **TU27**

FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C] Isotype Control

Conjugation

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM**

of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD122/IL-2RB Antibody and APC Anti-Human CD56 Antibody.

Preparation & Storage

Keep as concentrated solution. Storage

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names $CD122; IL-2RB; IL-2R\beta; Interleukin-2\ receptor\ subunit\ beta; p75$

Uniprot ID P14784 Gene ID 3560

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com Web:www.elabscience.com

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

Elabscience®

CD122 is a 70-75 kD type I transmembrane glycoprotein and member of the Ig superfamily. It is IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. CD122 is constitutively expressed by NK cells and at lower levels by a subset of T cells. Its expression is upregulated upon activation. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation.

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com