



A Reliable Research Partner in Life Science and Medicine

Recombinant GCSH Monoclonal Antibody

catalog number: AN300279P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human GCSH Protein


Host Rabbit Isotype lgG Clone B201 **Purification** Protein A


Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

1:100-1:500 IHC-P

Data





Immunohistochemistry of paraffin-embedded human kidney using GCSH Monoclonal Antibody at dilution of 1:200.

Immunohistochemistry of paraffin-embedded human liver using GCSH Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

Degradation of glycine is brought about by the glycine cleavage system, which is composed of four mitochondrial protein components: P protein (a pyridoxal phosphate-dependent glycine decarboxylase), H protein (a lipoic acidcontaining protein), T protein (a tetrahydrofolate-requiring enzyme), and L protein (a lipoamide dehydrogenase). The protein encoded by this gene is the H protein, which transfers the methylamine group of glycine from the P protein to the T protein. Defects in this gene are a cause of nonketotic hyperglycinemia (NKH). Two transcript variants, one protein-coding and the other probably not protein-coding, have been found for this gene. Also, several transcribed and non-transcribed pseudogenes of this gene exist throughout the genome.

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Rev. V1.2