FBXW8 Polyclonal Antibody

catalog number: E-AB-19858



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Description			
Reactivity	Human		
Immunogen	Synthetic peptide of human FBXW8		
Host	Rabbit		
Isotype	IgG		
Purification	Antigen affinity purification		
Conjugation	Unconjugated		
buffer	Phosphate buffered solution	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Applications Recommended Dilution		
IHC	1:40-1:200		
Data			
	ry of paraffin-embedded Human using FBXW8 Polyclonal Antibody th	Immunohistochemistry of paraffin-embedded Human ayroid cancer tissue using FBXW8 Polyclonal Antibody at	
at dilution of 1:70(×200) dilution of 1:70(×200)			
Preparation & Storage			
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.		
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the		
temperature recommended.			
Background			
This gene encodes a member of the F-box protein family, members of which are characterized by an approximately 40			

This gene encodes a member of the F-box protein family, members of which are characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into three classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene contains a WD-40 domain, in addition to an F-box motif, so it belongs to the Fbw class. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene.

For Research Use Only