



A Reliable Research Partner in Life Science and Medicine

IL17RA/CD217 Monoclonal Antibody

catalog number: AN200111P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human IL17RA protein


Host Mouse Isotype lgG1 Clone A959 **Purification** Protein A

Buffer 0.2 µm filtered solution in PBS

Applications Recommended Dilution

1:500-1:1000 **WB**

Data



Western Blot with IL17RA / CD217 Monoclonal Antibody at dilution of 1:500. Lane A: Hela Whole Cell Lysate,

Lysates/proteins at 30 µg per lane.

Observed-MW:105 kDa Calculated-MW:96 kDa

Preparation & Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of **Storage**

activity. Antibody products are stable for twelve months from date of receipt when

stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

Interleukin 17 (also known as CTLA-8) is a T cell-expressed pleotropic cytokine. IL-17 binds to IL-17 receptor A (IL-17 RA), also known as IL-17 receptor (IL-17 R). IL-17 RA shares no homology with any known family of receptors. While the expression of IL-17 is restricted to activated T cells, the IL-17 RA mRNA exhibits a broad tissue distribution, and has been detected in virtually all cells and tissues tested. Human IL-17 RA is a 120 kDa, 866 amino acid (aa) type I membrane glycoprotein with a 293 aa extracellular domain, a 21 aa carboxy-proximal transmembrane domain, and a 525 aa cytoplasmic tail. Within the ECD, human IL-17 RAshares 72% aa sequence identity with mouse and rat IL-17 RA. The signaling events of IL-17 includes activation of NF-kappa B and JNK, and require TNF receptor-associated factors 6 (TRAF6) in the signaling pathway.

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Rev. V1.1