

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Mouse CD16/32 Antibody[2.4G2]

Catalog Number: E-AB-F0997UC

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2b, κ
Clone No. 2.4G2

Isotype Control FITC Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843C]

Conjugation FITC

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

Applications Recommended usage


FCM Each lot of this antibody is quality control tested by flow cytometric analysis. Please

check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells

in 100 µL volume].

Data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD16/32 Antibody (filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD16a/b;CD32;CD32A/B;FCG2A;FCGR2A/BFCGR3;FCGR3A/B;Fc fragment of IgG low

affinity Illa/b receptor;Fc fragment of IgG low affinity Illb receptor;Fc fragment of IgG low

affinity lla/b receptor;Fc gamma Rlla/bFc gamma receptor III A/B;FcGR

Uniprot ID P08508;P08101

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID Background 14130,14131

CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and mediate adaptive immune responses.

Toll-free: 1-888-852-8623 Web:www.elabscience.com

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email:techsupport@elabscience.com

Rev. V1.5