



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD27 Antibody[HIBQ27-1]

catalog number: AN009420P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD27 protein

Host Mouse

 $\begin{tabular}{lll} \textbf{Isotype} & Mouse IgG1, \kappa \\ \end{tabular}$ $\begin{tabular}{lll} \textbf{Clone} & HIBQ27-1 \\ \end{tabular}$

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM 2 μ g/mL(0.5×10⁶-1×10⁶ cells)

Data



Verified Samples in FCM: Human peripheral blood lymphocytes

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.