# Purified Anti-Human CD27 Antibody[HIBQ27-1]

**catalog number:** **AN009420P**

**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human |
| **Immunogen** | Recombinant Human CD27 protein |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Clone** | HIBQ27-1 |
| **Purification** | >98%, Protein A/G purified |
| **Buffer** | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

| Applications | Recommended Dilution |
|---|---|
| **FCM** | ≤ 0.2 μg per million cells in 100 μL volume |

## Data




Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD27 Antibody[HIBQ27-1] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| **Shipping** | Ice bag |

## Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com