

GM-CSF/CSF2/MGI-I/MGI-IGM, Mouse, Recombinant

Cat. No. : PCK038

General Information

Synonyms	Granulocyte-macrophage colony-stimulating factor;Csf2;GM-CSF;Colony-stimulating factor;Csfgm
Species	Mouse
Expression host	E.coli
Sequence	Ala18-Lys141
Accession	P01587
Mol mass	14.2 KDa
Expiration date	12 months
Bio activity	Measured in a cell proliferation assay using FDC-P1 cells. The ED50 for this effect is 15-50 pg/mL.


Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 1 mM EDTA, 4% Mannitol, pH 8.0.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


Background

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) was initially characterized as a growth factor that can support the in vitro colony formation of granulocyte-macrophage progenitors. It is produced by a number of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cells and fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophage progenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. On mature hematopoietic, monocytes/ macrophages and eosinophils. GM-CSF has a functional role on nonhematopoietic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF can also stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma and adenocarcinoma cell lines.

SDS-PAGE



Bio activity



Procell[®]system

Procell[®]system