

GM-CSF/CSF2, Swine, Recombinant**Cat. No. : GPCK369****产品信息**

物种	Swine
表达宿主	E.coli
序列信息	APTRPPSPVTRPWQHVD AIKEALSLLNNSNDTAAVMNETVDVVCEMFDPQEPTCVQTRLNLY KQGLRGSLTRLKSPLTLLAKHYEQHCPLTEETSCETQSITFKSFKDSLNLKFLFTIPFDCWGPVK K with polyhistidine tag at the N-terminus.
检索号	Q29118.1
标签	His-tag at the N-terminus
分子量	15.3 kDa
有效期	12 months
生物活性	Measure by its ability to induce proliferation in TF-1 cells. The ED50 for this effect is < 3 ng/mL.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Granulocyte-macrophage colony-stimulating factor (GM-CSF), also known as colony-stimulating factor 2 (CSF2), is a monomeric glycoprotein secreted by macrophages, T cells, mast cells, natural killer cells, endothelial cells and fibroblasts that functions as a cytokine. GM-CSF also plays a role in embryonic development by functioning as an embryokine produced by reproductive tract.

