

Recombinant Mouse IL13RA1 Protein (His Tag)

Catalog Number: PKSM040873

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse IL13RA1 protein Met 1-Thr 340, with an C-terminal His
Calculated MW	37.6 kDa
Observed MW	50-55 kDa
Accession	NP_598751.3
Bio-activity	1. Measured by its binding ability in a functional ELISA. 2. Immobilized mouse IL13RA1-His at 10µg/mL (100µL/well) can bind human Fc-IL13, the EC ₅₀ of human Fc-IL13 is 0.5-1.5 µg/mL.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Interleukin 13 receptor, alpha 1, also known as IL13RA1/IL-13RA1 and CD213A1 (cluster of differentiation 213A1), is a subunit of the interleukin 13 receptor. This subunit forms a receptor complex with IL4 receptor alpha, a subunit shared by IL13 and IL4 receptors. IL13RA1/IL-13RA1 serves as a primary IL13-binding subunit of the IL13 receptor, and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2, and thus may mediate the signaling processes that lead to the activation of JAK1, STAT3 and STAT6 induced by IL13 and IL4. IL13RA1/IL-13RA1 binds with low affinity to interleukin-13 (IL13). This subunit together with IL4RA can form a functional receptor for IL13. IL13RA1/IL-13RA1 also serves as an alternate accessory protein to the common cytokine receptor gamma chain for interleukin-4 (IL4) signaling, but cannot replace the function of IL2RG in allowing enhanced interleukin-2 (IL2) binding activity.