

Recombinant Human Pancreasin/Marapsin/PRSS27 Protein (His Tag)

Catalog Number: PKSH030900

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Pancreasin/Marapsin/PRSS27 protein Met 1-Lys 290, with an C-terminal His
Calculated MW	30.9 kDa
Observed MW	42 kDa
Accession	Q9BQR3
Bio-activity	Measured by its ability to cleave a colorimetric peptide substrate, NcarbobenzoyloxyGlyArgThioBenzyl ester (ZGR-SBzl), in the presence of 5, 5'-Dithiobis (2-nitrobenzoic acid) (DTNB). The specific activity is > 2000 pmols/min/μg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.5

The name "Pancreasin" because it is transcribed strongly in the pancreas. This secreted, tryptic serine protease, also known as Marapsin or PRSS27 (Protease, serine, 27), which is a member of the peptidase S1 family. Pancreasin is inhibited by benzamidine and leupeptin but resists several classic inhibitors of trypsin. Marapsin was constitutively expressed in nonkeratinizing stratified squamous epithelia of human esophagus, tonsil, cervix, larynx, and cornea. In fact, marapsin was the second most strongly up-regulated protease in psoriatic lesions, where expression was localized to the upper region of the hyperplastic epidermis. Similarly, in the hyperproliferative epithelium of regenerating murine skin wounds, marapsin localized to the suprabasal layers, where keratinocytes undergo squamous differentiation. Marapsin's restricted expression, localization, and cytokine-inducible expression suggest a role in the terminal differentiation of keratinocytes in hyperproliferating squamous epithelia.