A Reliable Research Partner in Life Science and Medicine

## Purified Anti-Human CD163 Antibody[MAC2-158]

catalog number: AN004800P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human

**Immunogen** Recombinant Human CD163 protein

**Host** Mouse

IsotypeMouse IgGl,  $\kappa$ CloneMAC2-158

**Purification** >98%, Protein A/G purified


**Buffer** Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze

to completely remove the stabilizer prior to labeling.

# Applications Recommended Dilution

FCM  $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$ 

#### Data



Human peripheral blood monocytes cell were stained with 0.2  $\mu g$  Purified Anti-Human CD163 Antibody[MAC2-158] (Right) and 0.2  $\mu g$  Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

IgG Secondary Antibody.

### **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

#### Background

The protein encoded by this gene is a member of the scavenger receptor cysteine-rich (SRCR) superfamily, and is exclusively expressed in monocytes and macrophages. It functions as an acute phase-regulated receptor involved in the clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages, and may thereby protect tissues from free hemoglobin-mediated oxidative damage. This protein may also function as an innate immune sensor for bacteria and inducer of local inflammation. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.

#### For Research Use Only

Fax: 1-832-243-6017