

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 700 Anti-Human CD18 Antibody[60.3]

Catalog Number: E-AB-F1412M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human Host Mouse

**Isotype** Mouse IgG2a, κ

**Clone No.** 60.3

Isotype Control Elab Fluor®700 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802M1]

Conjugation Elab Fluor® 700

**Conjugation Information** Elab Fluor® 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

#### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### Data



Staining of normal human peripheral blood cells with Elab

Fluor<sup>®</sup> 700 Anti-Human CD18 Antibody[60.3](left) or Elab Fluor<sup>®</sup> 700 Mouse IgG2a, κ Isotype Control (right). Cells in the leucocyte gate were used for analysis.

#### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

#### **Antigen Information**

Alternate Names Integrin β2 subunit;LFA -1 β subunit;β2 integrin;lTGB2;b2 integrin;beta 2 integrin

Web: www.elabscience.cn

 Uniprot ID
 P05107

 Gene ID
 3689

### For Research Use Only



## **Elabscience Biotechnology Co., Ltd.**

A Reliable Research Partner in Life Science and Medicine

**Background** 

CD18, also known as integrin  $\beta2$  subunit, LFA-1  $\beta$  subunit, and  $\beta2$  integrin, is a 90 - 95 kD type I glycoprotein. CD18 non-covalently associates with CD11a, CD11b, or CD11c. CD18 is expressed on all leukocytes. CD18 and associated  $\alpha$  chains function in the adhesion and signaling in hematopoietic cells.