

A Reliable Research Partner in Life Science and Medicine

# Recombinant Rat CD62L/L-Selectin Protein(Fc Tag)

Catalog Number: PDMR100048

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Rat **Species** 

Source HEK293 Cells-derived Rat CD62L/L-Selectin proteins Trp39-Asn332, with an C-

**Calculated MW** 57.5 kDa Observed MW 75-85 kDa P30836 Accession

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 90% as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU/mg of the protein as determined by the LAL method

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% **Formulation** 

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Rat CD62L/L-Selectin proteins, 2 μg/lane of Recombinant Rat CD62L/L-Selectin proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 57.5 KD

## **Background**

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Rev. V1.8

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

L-Selectin (also known as Leukocyte Selectin, LAM-1, LECAM-1, LECCAM-1, TQ1, Leu-8, MEL-14 antigen, DREG, lymph node homing receptor, CD62L) is a member of the Selectin family of cell surface molecules which include E-Selectin and P-Selectin. All Selectins have an extracellular domain composed of an amino-terminal calciumdependent lectin domain, an epidermal growth factor (EGF)-like domain, two to nine short consensus repeat (SCR) units, a transmembrane domain, and a cytoplasmic tail. L-Selectin expression is limited to hematopoietic cells, with most leukocytes expressing L-Selectin at some stage of differentiation. The majority of myeloid cells, B cells, and virgin T cells express L-Selectin, while only a sub-population of memory T cells and NK cells express L-Selectin. Lymphocytes and neutrophils exhibit a reversible loss of L-Selectin after cellular activation that results from endoproteolytic release of the extracellular portion of receptor from the cell surface. Cleavage of L-Selectin from the cell surface results in a high circulating level of functionally active soluble L-Selectin. All selectins bind sialytated and fucosylated oligosaccharides that are linked to glycoproteins and glycolipids. L-Selectin specifically binds to at least three different heavily glycosolylated mucin-like proteins: GlyCAM-1, CD34, and MAdCAM-1. Multiple studies indicated that L-Selectin, P-Selectin E-Selectin collaborate to mediate the initial binding of leukocytes to endothelium at sites of tissue injury and inflammation, producing the characteristic "rolling" of leukocytes along the endothelium. L-Selectin knockout mice have a 70% decrease in rolling leukocytes in exposed mesentery and have impaired neutrophil and monocyte migration into areas of inflammation. Additionally, L-Selectin knockout mice have relatively few lymphocytes present in peripheral lymph nodes and Peyer's patches. Short-term in vivo homing experiments in L-Selectin deficient mice demonstrate that L-Selectin is involved in lymphocyte homing to Peyer's patches and mesenteric lymph nodes in the gut. Rat and human L-Selectin share 77% amino acid sequence homology. Rat and mouse L-Selection share 83% amino acid sequence homology.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com