

Recombinant Phospho-HER2/ErbB2 (Tyr1221, 1222) Monoclonal Antibody


catalog number: **AN300005L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human
Immunogen	A synthetic phosphopeptide corresponding to residues around (Tyr1221, 1222) of human HER2/ErbB2.
Host	Rabbit
Isotype	IgG
Clone	A1115
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS


Applications	Recommended Dilution
WB:	1:1000-1:5000

Data



Western blot analysis of extracts from serum-starved SkBr3, untreated (line A); treated with EGF (Cat: PKSH031641) (200 ng/mL, 10 min), without peptide (line B) or antigen-specific phosphopeptide (line C) or antigen-specific peptide (line D) using Phospho-HER2/ErbB2 (Tyr1221, 1222) Monoclonal Antibody at 1:5000 dilution.

Observed-MW:185 kDa
Calculated-MW:138 kDa



Western blot analysis of extracts from serum-starved SkBr3, untreated (line A); treated with EGF (Cat: PKSH031641) (200 ng/mL, 10 min) (line B); treated with EGF and λ-phosphatase (line C) using Phospho-HER2/ErbB2 (Tyr1221, 1222) Monoclonal Antibody at 1:5000 dilution.

Observed-MW:185 kDa
Calculated-MW:138 kDa

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq, Jul 2008]

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.5