A Reliable Research Partner in Life Science and Medicine

Recombinant Human FGF-2/FGF basic/FGFb Protein

Catalog Number: PKSH033503

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source E.coli-derived Human FGF-2/FGF basic/FGFb protein Met1-Ser155

 Mol_Mass
 17.2 kDa

 Accession
 BAG70135.1

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $\leq 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 200mM NaCl, pH 7.5.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Fibroblast growth factors (FGF) are a family of heparin-binding secreted proteins that stimulate cell proliferation and differentiation in a wide variety of tissues. FGFs play important roles in diverse biological functions both in vivo and in vitro; including mitogenesis; cellular migration; differentiation; angiogenesis; and wound healing. Human embryonic stem cell (hESC) cultures require FGF basic (also known as FGF-2 or bFGF) in cell culture media to remain in an undifferentiated and pluripotent state. Thermostable FGF basic was engineered for enhanced stability in culture media; without modification of its biological function. FGF basic is a required component of stem cell culture media for maintaining cells in an undifferentiated state. Because FGF basic is unstable; daily media changes are needed. The thermostable FGF basic that supports a 2-day media change schedule; so no media changes are required over a weeken d. This thermostable FGF basic was more stable than FGF basic in biochemical studies; and maintained cell growth; pluripotency and differentiation potential with a 2-day feeding schedule.

For Research Use Only

Fax: 1-832-243-6017