Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Human MMP-2 Antibody Pair Set

Catalog No. E-KAB-0451 Applications ELISA

Synonyms MMP2;CLG4;CLG4A;MMP-II;MONA;TBE-1;Gelatinase A

Kit components & Storage

Title	Specifications	Storage
Human MMP-2 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Human MMP-2 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0451)	
		Human MMP-2 Capture Antibody	Human MMP-2 Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human MMP-2 protien	Recombinant Human MMP-2 protien
Information	Swissprot	P08253	
Product details	Reactivity	Human	Human
	Host	Rabbit	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human MMP-2 in ELISAs.	

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn





A Reliable Research Partner in Life Science and Medicine

Applications

Human MMP-2 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human MMP-2 Capture	
Capture		Antibody	10
			ensity
ELISA	1:1000-1:10000	Human MMP-2 Detection	Optical Density
Detection		Antibody (Biotin)	•
			0.01
			0.1 10 100 Human MMP-2 Concentration(ng/mL)
			Transar (4.41 - 2 Concentration(ng/mz.)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes , such as embryonic development , reproduction , and tissue remodeling , as well as in disease processes , such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades type IV collagen , the major structural component of basement membranes. The enzyme plays a role in endometrial menstrual breakdown , regulation of vascularization and the inflammatory response.

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn