Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Porcine ANG1 Antibody Pair Set

Catalog No. E-KAB-0667 Applications ELISA

Synonyms ANG-1;ANGPT1;AGP1;AGPT

Kit components & Storage

Title	Specifications	Storage
Porcine ANG1 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze/thaw cycles.
Porcine ANG1 Detection Antibody	1 vial, 50 μL	Store at -20℃ for one year.
(Biotin)		Avoid freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0667)		
		Porcine ANG1 Capture Antibody	Porcine ANG1 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Porcine ANG1 protein	Recombinant Porcine ANG1 protein	
Information	Swissprot	Q9BDY8		
Product details	Reactivity	Porcine	Porcine	
	Host	Mouse	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5 mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%	
		glycerol; pH 7.5	protective protein; 50% glycerol; pH	
			7.5	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Porcine ANG1 in ELISAs.		

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn

Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

Applications

Porcine ANG1 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Porcine ANG1 Capture	
Capture		Antibody	10
			disa the state of
ELISA	1:1000-1:10000	Porcine ANG1 Detection	Optical Density
Detection		Antibody (Biotin)	5 0.1
			:
			0.01 100 1000 10000 Porcine ANG1 Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Angiopoietins are proteins with important roles in vascular development and angiogenesis. All angiopoietins bind with similar affinity to an endothelial cell-specific tyrosine-protein kinase receptor. The protein encoded by this gene is a secreted glycoprotein that activates the receptor by inducing its tyrosine phosphorylation. It plays a critical role in mediating reciprocal interactions between the endothelium and surrounding matrix and mesenchyme and inhibits endothelial permeability. The protein also contributes to blood vessel maturation and stability, and may be involved in early development of the heart. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Tel: 400-999-2100 Web: www.elabscience.cn Email: techsupport@elabscience.cn