

A Reliable Research Partner in Life Science and Medicine

Recombinant Purine nucleoside phosphorylase/PNP Monoclonal Antibody

catalog number: AN300415P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human Purine nucleoside phosphorylase/PNP protein


HostRabbitIsotypeIgGCloneB341PurificationProtein A

Buffer 0.2 μm filtered solution in PBS

Applications Recommended Dilution


WB 1:500-1:2000IP 1-4 μL/mg of lysate

Data



Western Blot with PNP3 Monoclonal Antibody at dilution of 1:500 dilution. Lane A: Jurkat Whole Cell Lysate, Lane B: K562 Whole Cell Lysate, Lane C: 293T Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:32 kDa Calculated-MW:32 kDa



Immunoprecipitation analysis using 2 μL anti-PNP-3
Monoclonal Antibody and 15 μl of 50 % Protein G agarose.
Western blot was performed from the immunoprecipitate using PNP-3 Monoclonal Antibody at a dilution of 1:100.
Lane A:0.5 mg Jurkat Whole Cell Lysate, Lane B:0.5 mg
K562 Whole Cell Lysate

Observed-MW:32 kDa Calculated-MW:32 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

This gene encodes an enzyme which reversibly catalyzes the phosphorolysis of purine nucleosides. The enzyme is trimeric, containing three identical subunits. Mutations which result in nucleoside phosphorylase deficiency result in defective T-cell (cell-mediated) immunity but can also affect B-cell immunity and antibody responses. Neurologic disorders may also be apparent in patients with immune defects. A known polymorphism at an position 51 that does not affect enzyme activity has been described. A pseudogene has been identified on chromosome 2.

For Research Use Only