

# IL-32 Polyclonal Antibody(Capture)

catalog number: AN000680P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Human
<b>Immunogen</b>	Recombinant Human IL-32 protein expressed by E.coli
<b>Host</b>	Rabbit
<b>Isotype</b>	Rabbit IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
----------------------	-----------

## Data



Sandwich ELISA-Recombinant Human IL-32 protein standard curve. Background subtracted standard curve using IL-32 antibody (AN000680P) (Capture), IL-32 Antibody (AN001830P) (Detector) in sandwich ELISA. The reference range value for Recombinant Human IL-32 protein is 15.63-1000 pg/mL.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

IL-32 is a recently discovered cytokine that induces various proinflammatory cytokines (TNF-alpha, IL-1beta, IL-6) and chemokines in both human and mouse cells through the NF-kappaB and p38 MAPK inflammatory signal pathways. It is regulated robustly by other major proinflammatory cytokines and is crucial to inflammation and immune responses. Four of the IL-32 isoforms (alpha, beta, gamma, and delta) are the most representative IL-32 transcripts, and the gamma isoform of IL-32 is the most active, although all isoforms are biologically active. IL-32, a cytokine produced mainly by T, natural killer, and epithelial cells induces significant amounts of TNFalpha and MIP-2 and increases the production of both cytokines in a dose-dependent manner. IL-32 has been implicated in inflammatory disorders, Mycobacterium tuberculosis infections, inflammatory bowel disease, and influenza A virus infection, as well as in some autoimmune diseases, such as rheumatoid arthritis, ulcerative colitis, and in the human stomach cancer, human lung cancer, and breast cancer tissues. Thus, IL-32 expression might be valuable as a biomarker for cancer.

## For Research Use Only