

Recombinant Bovine Annexin V Protein (His Tag)

Catalog Number: PDEB100007

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Bovine
Source	E.coli-derived Bovine Annexin V protein Met1-Asp320, with an N-terminal His
Calculated MW	35.1 kDa
Observed MW	35 kDa
Accession	P81287
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Bovine Annexin V proteins, 2 µg/lane of Recombinant Bovine Annexin V proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 35 kDa.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Peptidyl-prolyl cis-trans isomerase A, also known as PPIase A, Rotamase A, Cyclophilin A, Cyclosporin A-binding protein, PPIA and CYPA, is a cytoplasm protein which belongs to the cyclophilin-type PPIase family and PPIase A subfamily. Cyclophilins (CyPs) are a family of proteins found in organisms ranging from prokaryotes to humans. These molecules exhibit peptidyl-prolyl isomerase activity, suggesting that they influence the conformation of proteins in cells. PPIA / Cyclophilin A accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. PPIA / Cyclophilin A is secreted by vascular smooth muscle cells in response to inflammatory stimuli, and could thus contribute to atherosclerosis. It is not essential for mammalian cell viability. PPIA / Cyclophilin A can interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions.