Purified Anti-Human CD200R1 Antibody[OX-108]

catalog number: E-AB-F14750P




Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD200R1 protein |
| Host | Mouse |
| Isotype | Mouse IgG1,ĸ |
| Clone | OX-108 |
| Purification | >98%, Protein A/G purified |
| Conjugation | Unconjugated |
| buffer | PBS, pH 7.2. Contains 0.05% proclin 300. |
| | |

ApplicationsRecommended DilutionFCM2 μg/mL(1×10⁵-5×10⁵ cells)

Data



Human peripheral blood lymphocytes were stained with 0.2

μg Purified Anti-Human CD200R1 Antibody[OX-108]

(Right) and 0.2 µg Mouse IgG1, κ Isotype Control (Left),

followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

IgG Secondary Antibody, then anti-Human CD4 PerCP-

conjugated Monoclonal Antibody.

| Preparation & Storage | |
|-----------------------|---|
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
| | thaw cycles. |
| Shipping | Order now, ship in 3 days |
| | |

Background

CD200R, also known as OX2R, is a membrane glycoprotein with up to 70% of its weight derived from N-linked glycosylation. CD200R is expressed primarily by monocytes and neutrophils but also by other leukocytes including T lymphocytes and mast cells. The interaction between CD200 and CD200R may contribute to pathways that suppress and limit macrophage induced inflammatory damage in tissue. Studies suggest that CD200-CD200R interaction may be involved in the control of myeloid cellular function. Levels of expression on resting peripheral blood cells are relatively low.

For Research Use Only