

A Reliable Research Partner in Life Science and Medicine

PE/Cyanine 5 Anti-Human CD138/Syndecan-1 Antibody[B-B4]

Catalog Number: E-AB-F1411G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ

Clone No. B-B4

Isotype Control PE/Cyanine5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792G]

Conjugation PE/Cyanine 5

Conjugation Information PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Staining of the U266 cell line with PE/Cyanine5 Anti-Human CD138/Syndecan-1 Antibody[B-B4] (filled gray histogram) or PE/Cyanine5 Mouse IgG1, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Web: www.elabscience.cn

Shipping lce bag

Antigen Information

Alternate Names Syndecan-1
Uniprot ID P18827
Gene ID 6382

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Background

CD138, a member of the syndecan protein family, is a type I integral membrane heparin sulfate proteoglycan also known as Syndecan-1. Syndecan-1 participates in cell proliferation, cell migration, and cell-matrix adhesion via interaction with collagen, fibronectin, and other soluble molecules (such as FGF-basic). It is expressed on normal and malignant human plasma cells, pre-B cells, epithelial cells, and endothelial cells.