A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse Thrombomodulin/THBD Protein (His Tag)

Catalog Number: PKSM041274

Note: Centrifuge before opening to ensure complete recovery of vial contents.

-			
- 1	00	crip	IOI
JU			

Species Mouse

Source HEK293 Cells-derived Mouse Thrombomodulin/THBD protein Leu17-Ser517, with an

C-terminal His

 Calculated MW
 54.6 kDa

 Observed MW
 85-90 kDa

 Accession
 P15306

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Concentration Subject to label value.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.


Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.

Data



Background

Thrombomodulin is also known as CD141 antigen and blood dendritic cell antigen 3 (BDCA3), which is encoded by the THBD gene. The deduced amino acid sequence of mouse THBD predicts a signal peptide (aa 1 to 16) and a mature chain (aa 17 to 577) that consists of the following domains: C-type lectin, EGF-like, transmembrane and cytoplasmic. Mouse THBD is corresponding to the extracellular portion of the type I membrane protein. Predominantly synthesized by vascular endothelial cells, THBD inhibits coagulation and fibrinolysis. It functions as a cell surface receptor and an essential cofactor for active thrombin, which in turn activates protein C and thrombinactivatable fibrinolysis inhibitor (TAFI), also known as carboxypeptidase B2 (CPB2). In addition, THBD gene polymorphisims are associated with human disease and THBD plays a role in thrombosis, stroke, arteriosclerosis, and cancer.