

Recombinant Human SETD7/SET7/9 Protein (His Tag)

Catalog Number: PKSH031208

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human SETD7/SET7/9 protein Asp 2-Lys 366, with an N-terminal His
Calculated MW	41.5 kDa
Observed MW	48 kDa
Accession	NP_085151.1
Bio-activity	Not validated for activity

Properties

Purity	> 94 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual.

Data



> 94 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Histone-lysine N-methyltransferase SETD7, also known as SET domain containing (lysine methyltransferase) 7, SET7/9, Histone H3-K4 methyltransferase SETD7, H3-K4-HMTase SETD7, and SETD7, is a member of the histone-lysine methyltransferase family and SET7 subfamily. SETD7 is widely expressed and expressed in pancreatic islets. SETD7 contains three MORN repeats and one SET domain. SETD7 plays a central role in the transcriptional activation of genes such as collagenase or insulin. As a protein lysine methyltransferase (PKMT), SETD7 also has methyltransferase activity toward non-histone proteins such as p53/TP53, TAF10, and possibly TAF7 by recognizing and binding in substrate proteins. The mono-methyltransferase activity of SETD7 is achieved by disrupting the formation at near-attack conformations for the dimethylation reaction. SETD7 is also a novel coactivator of NF-kappaB and plays a role in inflammation and diabetes.