IL-17 RA/IL-17 R Polyclonal Antibody(Capture/Detector)

catalog number: AN000470P



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-						٠,		
D	Α		C	PТ	m	ŤΤ	U.	m
\mathbf{L}	v	0	v.	ш	עו	UΙ	U.	

Reactivity Human

Immunogen Recombinant Human IL-17 RA/IL-17 R RA protein expressed by Mammalian

Host Rabbit
Isotype Rabbit IgG

Purification Antigen Affinity Purification


Conjugation Unconjugated

buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications Recommended Dilution

ELISA Capture 2-8 μg/mL **ELISA Detector** 0.1-0.4 μg/mL

Data



Sandwich ELISA-Recombinant Human IL-17 RA/IL-17 R

RA protein standard curve.Background subtracted standard

curve using IL-17 RA/IL-17 R antibody(AN000470P)

(Capture), IL-17 RA/IL-17 R antibody (AN000470P)

(Detector) in sandwich ELISA. The reference range value for

Recombinant Human IL-17 RA/IL-17 R RA protein is 0.31-

20 pg/mL.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different isoforms.

For Research Use Only