

FGF-2/bFGF/FGF-b, Rat, Recombinant**Cat. No. : PCK012****产品信息**

别名	Fibroblast Growth Factor 2;FGF-2;Basic Fibroblast Growth Factor;bFGF;Heparin-Binding Growth Factor 2;HBGF-2;FGF2;FGFB
物种	Rat
表达宿主	E.coli
序列信息	Ala11-Ser154
检索号	P13109
分子量	16.2 kDa
有效期	12 months
生物活性	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 0.3-1.8 ng/mL.

产品特性

内毒素	< 1.0 EU per 1 µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation, growth, and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis, cell growth, pattern formation, embryonic development, metabolic regulation, cell migration, neurotrophic effects, and tissue repair. FGF-basic is a non-glycosylated heparin binding Growth Factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4.




SDS-PAGE

普诺赛®



Bio activity



普诺赛® | Procell system

普诺赛® | Procell system

