A Reliable Research Partner in Life Science and Medicine

Recombinant Human 5'-Nucleotidase/5'-NT/CD73 (P21589, C-6His)

Catalog Number: PKSH033860

Note: Centrifuge before opening to ensure complete recovery of vial contents.

-			
- 1	00	crip	IOI
JU			

Species Human

Source HEK293 Cells-derived Human 5'-Nucleotidase;5'-NT;CD73 protein Trp27-Lys547, with

an C-terminal His

Calculated MW58.8 kDaObserved MW62 kDaAccessionP21589

Bio-activity Immobilized Human CD73-His at 2μg/ml (100 μl/well) can bind Anti-Human CD73

mAb. The ED₅₀ of Anti-Human CD73 mAb is 2.78 ng/ml.

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Concentration Subject to label value.

Endotoxin $\leq 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ **Storage** Storage Stor


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 120mM NaCl, 4mM CaCl₂,

20% Glycerol, pH 7.5.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD73 is an ecto 5'Nucleotidase expressed by most cell types. CD73 hydrolyzes extracellular nucleotides into membrane permeable nucleosides. CD73 is one of several enzymes responsible for the production of extracellular adenosine, a signaling molecule that is involved in responses to inflammation and tissue injury. CD73 is a lymphocyte maturation marker that has functions independent of its catalytic activity. CD73 is also a regulator of leukocyte extravasation, a function that requires its 5'Nucleotidase activity. Defects in NT5E are the cause of calcification of joints and arteries (CAJA). The recombinant CD73 lacking GPI anchor is secreted as a monomer.